



University
of Glasgow

Qin, S., Cao, Y., Luo, Y., Jiang, S., Clark, J. S., Wang, X. and Yang, G. (2019) Multi-gram scale synthesis of chiral 3-methyl-2,5-trans-tetrahydrofurans. *Helvetica Chimica Acta*, 102(7), e1900131.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

This is the peer reviewed version of the following article Qin, S., Cao, Y., Luo, Y., Jiang, S., Clark, J. S. , Wang, X. and Yang, G. (2019) Multi-gram scale synthesis of chiral 3-methyl-2,5-trans-tetrahydrofurans. *Helvetica Chimica Acta*, 102(7), e1900131, which has been published in final form at <http://dx.doi.org/10.1002/hlca.201900131>. This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

<http://eprints.gla.ac.uk/190218/>

Deposited on: 12 July 2019

Multi-Gram Scale Synthesis of Chiral 3-Methyl-2,5-*trans*-tetrahydrofurans

Shuanglin Qin,^{a,b} Yuting Cao,^b Yunhao Luo,^b Shende Jiang,^a J. Stephen Clark,^c Xiaoji Wang,^{*d,e}

Guang Yang^{*b}

^a School of Pharmaceutical Science and Technology, Tianjin University, Tianjin 300072, People's Republic of China.

^b The State Key Laboratory of Medicinal Chemical Biology, College of Pharmacy, Nankai University, Tianjin 300071, People's Republic of China, e-mail: Guang.yang@nankai.edu.cn

^c WestCHEM, School of Chemistry, Joseph Black Building, University of Glasgow, University Avenue, Glasgow G12 8QQ, United Kingdom

^d Engineering Research Center of Health Food Design & Nutrition Regulation, School of Chemical Engineering and Energy Technology, Dongguan University of Technology.

^e School of Life Science, Jiangxi Science and Technology Normal University, Nanchang, China. e-mail: 2012207455@tju.edu.cn.

In this paper, we report the rapid and facile synthesis of chiral 3-methyl-2,5-*trans*-tetrahydrofurans. This reaction utilizes cheap and easily available starting materials. A domino hydrolysis and intramolecular Michael-type ring closure reaction was the key step. As a result, synthesis of the desired 3-methyl-2,5-*trans*-tetrahydrofurans could be achieved in gram-scale over 7 linear steps with high chemical yield and high diastereoselectivity.

Keywords: 3-methyl-2,5-*trans*-tetrahydrofurans; Ring closure; Gram-scale synthesis; Highly diastereoselective.

Introduction

3-Methyl-2,5-*trans*-tetrahydrofurans are critical structural elements for various natural products, biologically active compounds, and pharmaceuticals. ^[1-7] For instance, natural products chagosensine, ^[8, 9] cationomycin, ^[10, 11] amphidinolides C, C2, C3 and F ^[12-15] possesses such *trans*-tetrahydrofuran substructures. (Figure 1) As a consequence, many methodologies have been developed for the construction of this type of heterocycle. ^[16-22]

Among them, intramolecular conjugated addition of an oxygen nucleophile is often used for the formation of 3-methyl-2,5-*trans*-tetrahydrofurans (Scheme 1a), which inspired us to identify suitable conditions to obtain useful amount of 3-methyl-2,5-*trans*-tetrahydrofurans intermediates for further studies on total synthesis of natural products. ^[23, 24] Herein, we reported a rapid and facile process to access gram-scale quantities of 3-methyl-2,5-*trans*-tetrahydrofurans in just 7 linear steps. A domino hydrolysis and diastereoselective intramolecular Michael-type ring closure strategy is the key step (Scheme 1b).






Figure 1. Representative natural products and bioactive compounds containing a 3-methyl-2,5-*trans*-tetrahydrofurans core.

a. previous works (Roush's work(1) and Spilling's work(2))



b. this work (domino hydrolysis and intramolecular Michael-type ring closure)




Scheme 1. Synthesis of 3-methyl-2,5-*trans*-tetrahydrofurans.

Results and Discussion


As depicted in Scheme 2, our synthetic route began with diazotization of inexpensive starting material, *L*-glutamic acid **1**, using NaNO_2 in HCl aqueous solution to provide lactone **2**. Reduction of the carboxyl group of lactone **2** with borane followed by protection of the primary alcohol **3** with TBSCl led to lactone **4**. Selective methylation of **4** using LiHMDS and MeI delivered **5** as a mixture of two diastereoisomers ($dr = 10:1$). Although the two stereoisomers of **5** could be separated by chromatography on silica gel, purification would be very time-consuming due to the similar polarity

of the two isomers. Hence, we used the mixture directly for the next step, and easy separation of isomeric products was performed at a later stage. With the mixture of two stereoisomers of **5** in hand, DIBAL reduction followed by Wittig reaction produced the pure conjugated ester **6** with an isolated yield over 85% of two steps. Subsequent Mitsunobu reaction of hydroxyl group in **6** delivered PNB ester **7**.



Scheme 2. The synthesis of intermediate **7**. *Reagents and conditions:* **a** NaNO₂, HCl, H₂O, 0 °C to rt; **b** BH₃·SMe₂, THF, 0 °C; **c** TBSCl, imidazole, DMAP, CH₂Cl₂, rt, 42% over 3 steps; **d** LiHMDS, MeI, THF, –78 °C, 80%; **e** DIBAL, –78 °C; **f** (Ph₃P)₃PCHCO₂Et, PhMe, 80 °C, 85% over 2 steps; **g** Ph₃P, DIAD, *p*-nitrobenzoic acid, THF, 0 °C, 74% over 3 steps.

Following successful preparation of ester precursor **7**, the 3-methyl-2,5-*trans*-tetrahydrofuran **8** was formed through a domino hydrolysis and intramolecular Michael-type ring closure process using K₂CO₃ as base with multi-gram scale (7.9 g). Notably, the methyl-bearing stereogenic centre played a very important role in controlling the stereochemical outcome of the ring closure reaction (Scheme 3). Under the same conditions, the ester **10** was obtained from the cyclisation of **9**, which was synthesised from lactone **4** using the same synthetic route as described in scheme 2, but with no diastereocontrol (*dr* = 1:1).



Scheme 3. Domino hydrolysis and intramolecular Michael-type ring closure reaction. *Reagents and conditions:* **a** K₂CO₃, EtOH, 55 °C, 89%; **b** DIBAL, –78 °C; **c** (Ph₃P)₃PCHCO₂Et, PhMe, 80 °C; **d** K₂CO₃, EtOH, 55 °C.

Conclusions

In summary, we have developed a domino hydrolysis and intramolecular Michael-type ring closure strategy to prepare 3-methyl-2,5-*trans*-tetrahydrofurans. It uses the readily-available chiral pool material L-glutamic acid as starting material. The required 3-methyl-2,5-*trans*-tetrahydrofuran was obtained with high chemical yield (18.8% overall yield) and high diastereoselectivity (*dr* > 25:1) over 7 steps. The process was also efficiently realized on multi-gram scale. Furthermore, the *trans*-tetrahydrofurans intermediate **8** can be subjected to further chain extension in either direction, which

might be useful for the synthesis of various natural products and biologically active compounds ^[25–29].

Supplementary Material

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/MS-number>.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (NSFC; No. 81703343 and 81503019), Fundamental Research Funds for the Central Universities. Tianjin Science and Technology Project [No. 17PTSYJC00070].

Author Contribution Statement

S. Qin and Y. Cao contributed equally to this work. S. Qin, Y. Cao, and Y. Luo performed the experiments, analyzed the data. Prof. S. Jiang and Prof. J. S. Clark gave important suggestions for this project and help improved the language. Prof. X. Wang and Dr. G. Yang designed the experiments and wrote the manuscript.

References

- [1] K. Kong, D. Romo, C. Lee, 'Enantioselective Total Synthesis of the Marine Toxin (–)-Gymnodimine Employing a Barbier-Type Macrocyclization', *Angew. Chem., Int. Ed.* **2009**, *48*, 7402–7405.
- [2] K. Kong, Z. Moussa, C. Lee, D. Romo, 'Total Synthesis of the Spirocyclic Imine Marine Toxin (–)-Gymnodimine and an Unnatural C4-Epimer', *J. Am. Chem. Soc.* **2011**, *133*, 19844–19856.
- [3] M. Litaudon, J. Hart, J. Blunt, R. Lake, M. Munro, 'Isohomohalichondrin B, a new antitumour polyether macrolide from the New Zealand deep-water sponge *Lissodendoryx* sp.', *Tetrahedron lett.* **2010**, *35*, 9435–9438.
- [4] M. Nakamura, T. Ohno, S. Kunitomo, H. Naganawa, T. Takeuchi, 'Kijimicin: An inhibitor of Human Immunodeficiency Virus in Acutely and Chronically Infected Cells', *J. Antibiot.* **1991**, *44*, 569–571.
- [5] L. Steinrauf, M. Pinkerton, J. Chamberlin, 'The structure of nigericin', *Biochem. Biophys. Res. Co.* **1968**, *33*, 29–31.
- [6] A. Mai, K. Kumagai, M. Tsuda, Y. Konishi, A. Tominaga, M. Tsuda, E. Fukushima, J. Kawabata, 'Iriomoteolide-13a, a cytotoxic 22-membered macrolide from a marine dinoflagellate *Amphidinium* species', *Tetrahedron.* **2014**, *70*, 2962–2965.
- [7] V. Havlíček, M. Ryska, S. Pospíšil, 'Negative-ion fast atom bombardment tandem mass spectrometry of sodium salts of monensins and related compounds', *J. Mass. Spectrom.* **1995**, *30*, 1089–1094.
- [8] M. Heinrich, J. Murphy, M. Ilg, A. Letort, J. Flasz, P. Philipps, A. Fuerstner, 'Total Synthesis of Putative Chagosensine', *Angew. Chem., Int. Ed.* **2018**, *57*, 13575–13581.
- [9] T. Rezanka, L. Hanus, V. Dembitsky, 'A Simple Synthetic Access to Differently 4-Substituted Neu5Ac2en Glycals Combining Elements of Molecules with Anti-Neuraminidase Activity', *Eur. J. Org. Chem.* **2003**, 4065–4077.
- [10] G. Nakamura, K. Kobayashi, T. Sakurai, K. Isono, 'Cationomycin, A New Polyether Ionophore Antibiotic Produced by *Actinomadura* nov. sp.', *J. Antibiot.* **1981**, *34*, 1513–1514.
- [11] G. Nakamura, K. Isono, 'A New Species of *Actinomadura* Producing A Polyether Antibiotic, Cationomycin', *J. Antibiot.* **1983**, *36*, 1468–1472.
- [12] J. Kobayashi, 'Amphidinolides and Its Related Macrolides from Marine Dinoflagellates', *J. Antibiot.* **2008**, *61*, 271–284.
- [13] J. Kobayashi, M. Ishibashi, 'Bioactive metabolites of symbiotic marine microorganisms', *Chem. Rev.* **1993**, *93*, 1753–1769.
- [14] J. Kobayashi, M. Tsuda, 'Amphidinolides, bioactive macrolides from symbiotic marine dinoflagellates', *Nat. Pro. Rep.* **2004**, *21*, 77–93.
- [15] J. Kobayashi, S. Kubota, M. Tsuda, 'Bioactive macrolides and polyketides from marine dinoflagellates', *Pure Appl. Chem.* **2003**, *75*, 337–342.
- [16] G. Valot, C. Regens, D. O'Malley, E. Godineau, H. Takikawa, A. Fürstner, 'Total Synthesis of Amphidinolide F', *Angew. Chem., Int. Ed.* **2013**, *52*, 9534–9538.

- [17] T. Aicher, K. Buszek, F. Fang, C. Forsyth, S. Jung, Y. Kishi, M. Matelich, P. Scola, D. Spero, S. Yoon, 'Total synthesis of halichondrin B and norhalichondrin B', *J. Am. Chem. Soc.* **1992**, *114*, 3162–3164.
- [18] G. Valot, D. Mailhol, C. Regens. D. O'Malley, E. Godineau, H. Takikawa, P. Philipps, A. Fürstner, 'Concise Total Syntheses of Amphidinolides C and F', *Chem-Eur. J.* **2015**, *21*, 2398–2408.
- [19] D. Mohapatra, P. Dasari, H. Rahaman, R. Pal, 'Stereoselective synthesis of the densely functionalized C1–C9 fragment of amphidinolides C and F', *Tetrahedron Lett.* **2009**, *50*, 6276–6279.
- [20] C. Song, H. Liu, M. Hong, Y. Liu, F. Jia, L. Sun, Z. Pan, J. Chang, 'Convergent Formal Synthesis of (±)-Roseophilin', *J. Org. Chem.* **2012**, *77*, 704–706.
- [21] Y. Liu, Y. Guo, F. Ji, D. Gao, C. Song, J. Chang, 'Divergent Syntheses of Carbazole Alkaloids Clausenapin, Indizoline, Claulansine M, and Clausenaline D', *J. Org. Chem.* **2016**, *81*, 4310–4315.
- [22] T. Kubota, M. Tsuda, J. Kobayashi, 'Absolute Stereochemistry of Amphidinolide C', *Org. Lett.* **2001**, *3*, 1363–1366.
- [23] R. H. Bates; J. B. Shotwell, W. Roush, 'Stereoselective Syntheses of the C(1)–C(9) Fragment of Amphidinolide C', *Org. Lett.* **2008**, *10*, 4343–4346.
- [24] M. P. Paudyal, N. P. Rath, C. D. Spilling, 'A Formal Synthesis of the C1–C9 Fragment of Amphidinolide C Employing the Tamaru Reaction', *Org. Lett.* **2010**, *12*, 2954–2957.
- [25] J. S. Clark, G. Yang, A. P. Osnowski, 'Synthesis of the C-1–C-17 Fragment of Amphidinolides C, C2, C3, and F', *Org. Lett.* **2013**, *15*, 1460–1463.
- [26] J. S. Clark, G. Yang, A. P. Osnowski, 'Synthesis of the C-18–C-34 Fragment of Amphidinolides C, C2, and C3', *Org. Lett.* **2013**, *15*, 1464–1467.
- [27] S. Mahapatra, R. G. Carter, 'Exploiting Hidden Symmetry in Natural Products: Total Syntheses of Amphidinolides C and F', *J. Am. Chem. Soc.* **2013**, *135*, 10792–10803.
- [28] J. Wang, J. Wang, C. Li, Y. Meng, J. Wu, C. Song, J. Chang, 'Synthesis of 5-*epi*-Taiwaniaquinone G', *J. Org. Chem.* **2014**, *79*, 6354–6359.
- [29] S. Mahapatra, R. G. Carter, 'Enantioselective Total Synthesis of Amphidinolide F, *Angew. Chem., Int. Ed.* **2012**, *51*, 7948–7951.